

Invia® SilverIon® NPWT

ANTIMICROBIAL WOUND CONTACT DRESSING

Effective

 The patented technology unfolds its power very quickly with continuous release of high concentrations silver ions in situ during wear time.

Simple

 Can be cut or overlapped to fit any wound size.

Comfortable

 Non-adherent for comfortable dressing changes ¹.



Providing choices for Negative Pressure Wound Therapy

Invia Silverlon NPWT Antimicrobial Wound Contact Dressing is intended to assist with the removal of fluids, exudates, infectious materials, and deliver antimicrobial silver ions in the dressing when activated by moisture. The silver ions in the dressing kill wound bacteria held in the dressing and provide an antimicrobial barrier for bacterial penetration of the dressing which may help reduce infection.

Invia Silverlon NPWT Antimicrobial Wound Contact Dressing is intended to be used in conjunction with Invia negative pressure wound therapy (NPWT), for local management of:


- 1st and 2nd degree burns
- Donor and graft site wounds
- Traumatic and surgical wounds

- Diabetic, pressure and venous ulcers
- Infected wounds*

Invia Silverlon NPWT Antimicrobial Wound Contact Dressing is a single layer of knitted nylon fiber substrate coated with metallic silver.

This Antimicrobial Wound Contact Dressing model is knitted to a specified pattern designed to allow optimal flow rate of wound exudates when used as the primary wound contact layer in a NPWT dressing set.

It is composed of 100% multifilament 15 denier nylon fiber, which is circumferentially coated with metallic silver at the rate of 0.68 mg/cm² (6.8 g/m²).



lonic silver is a broad spectrum antimicrobial, which has been shown to be effective against a range of pathogens associated with wound infection. Invia-Silverlon NPWT dressings have been tested ² in vitro and found effective against microorganisms such as:

- Gram Positive Bacteria: Staphylococcus aureus, Enterococcus faecalis, Staphylococcus epidermidis

Local contact

Gram Negative Bacteria: Pseudomonas aeruginosa, Enterobacter cloacae, Klebsiella pneumoniae and Acinetobacter
 Baumannii

Invia Silverlon NPWT Wound Contact Dressings are indicated for a total duration of contact (i.e. where treatment involves consecutive application of individual dressings) of up to 30 days.

Medical Vacuum Technology for Healthcare Professionals

Please contact us or your local Medela representative for details.

Medela and Invia are trademarks of Medela Holding AG, Switzerland. Silverlon is a trademark of Argentum Medical AG, USA.

Argentum Medical LLC 2571 Kaneville Court, Geneva, Illinois 60134 USA

Tel. +1 630 232 2507 Fax. +1 630 232 8005 EC REP

Emergo Europe Molenstraat 15 2513 BH The Hague The Netherlands Tel. +31 70 345 8570

United Kinadom

Medela UK Ltd.
Huntsman Drive, Northbank Industrial Park
Irlam, Manchester M44 5EG
United Kingdom
Phone +44 161 776 0400
Fax +44 161 776 0444
info@medela.co.uk
www.medela.co.uk

International Sales

Medela AG Lättichstrasse 4b 6341 Baar Switzerland Phone +41 41 562 51 51 Fax +41 41 562 51 00 ism@medela.ch www.medela.com

^{*} Invia-Silverlon NPWT Wound Contact Dressings may be used on those infected wound types listed above. Where used on infected wounds, the infection should be treated as per local clinical protocol.

¹ H. J. Siegel, D. F. Herrera, and J. Gay, "Silver negative pressure dressing with vacuum-assisted closure of massive pelvic and extremity wounds" Clinical Orthopaedics and Related ResearchR, vol. 472, no. 3, pp. 830–835, jun 2013.

² Data on file at Argentum Medical.